

PP - WAVES

The original article: Physical Review Letters **3**, 571 (1959)

SOME GRAVITATIONAL WAVES*

Asher Peres

Department of Physics, Israel Institute of Technology, Haifa, Israel

(Received October 19, 1959)

The few hitherto known wave-like exact solutions of the Einstein gravitational equations represented either plane^{1,2} or cylindrical³ waves. We here intend to derive a new class of solutions, displaying a lesser degree of symmetry, and thus more generality.

Let us consider the metric

$$ds^2 = -dx^2 - dy^2 - dz^2 + dt^2 - 2f(x, y, z + t)(dz + dt)^2.$$

One can easily show that $R_{zz} = R_{zt} = R_{tt} = f_{xx} + f_{yy}$, and other components vanish. If f is a harmonic function of x and y , this metric thus satisfies Einstein's equations in vacuo, whatever may be its dependence on $(z + t)$.

A convenient tetrad of orthonormal vectors is

$$h_{(1)}^k = (\cos a, \sin a, 0, 0),$$

$$h_{(2)}^k = (-\sin a, \cos a, 0, 0),$$

$$h_{(3)}^k = (0, 0, 1 - f, f),$$

$$h_{(4)}^k = (0, 0, -f, 1 + f),$$

where $\tan(2a) = (f_{xy}/f_{xx})$. The only nonvanishing independent physical components⁴ of the curvature tensor are

$$\sigma = R_{(m1n1)} = -R_{(m2n2)} = (f_{xx}^2 + f_{xy}^2)^{1/2},$$

where m and n take the values 3 and 4 only. Our metric thus belongs to the second class of Petrov's classification.⁴

Let us examine two examples. If

$$f = (x^2 - y^2) \sin(z + t),$$

one has $\sigma = 2 \sin(z + t)$. This is a plane monochromatic wave.

Another example is

$$f = xy(x^2 + y^2)^{-2} \exp[b^2 - (z + t)^2]^{-2} \quad \text{for } |z + t| < b,$$

$$f = 0 \quad \text{for } |z + t| \geq b.$$

(Note that f has derivatives of all orders at $|z + t| = b$.) This is a wave packet travelling with unit velocity in the negative z direction. It contains the singular segment $x = y = 0$, $-b - t < z < b - t$. The question of the existence of stable regular gravitational wave packets is still open.

If $f_{xx} + f_{yy} \neq 0$, one still has $g^{mn}R_{mn} = 0$ and $g^{mn}R_{mr}R_{ns} = 0$. The source of f can then be interpreted as a null electromagnetic field.⁵

*Partly supported by the U. S. Air Force, through the Air Research and Development Command.

¹H. Bondi, Nature **179**, 1072 (1957); H. Bondi, F. A. E. Pirani, and I. Robinson, Proc. Roy. Soc. (London) A **251**, 519 (1959).

²H. Takeno, Tensor **7**, 97 (1957); **8**, 59 (1958), and **9**, 76 (1959).

³N. Rosen, Bull. Research Council Israel **3**, 328 (1954).

⁴F. A. E. Pirani, Phys. Rev. **105**, 1089 (1957).

⁵C. W. Misner and J. A. Wheeler, Ann. Phys. **2**, 525 (1957).